arXiv:1211.2598v1 [cond-mat.stat-mech] 12 Nov 2012

ZUBAREV NONEQUILIBRIUM STATISTICAL OPERATOR IN RENYI STATISTICS. REACTION-DIFFUSION PROCESSES

P. Kostrobij and R. Tokarchuk
National University "Lvivs'ka Politekhnika", 79013 Lviv, 12 Bandera Str.

M. Tokarchuk*
Institute for Condensed Matter Physics NAS of Ukraine,
79011 Lviv, 1 Svientsitskii Str. and
National University "Lvivs'ka Politekhnika", 79013 Lviv, 12 Bandera Str.

B. Markiv
Institute for Condensed Matter Physics NAS of Ukraine,
79011 Lviv, 1 Svientsitskii Str.

(Dated: November 13, 2012)

Abstract

A statistical approach for consistent description of reaction-diffusion processes in system "gas-adsorbate-metal" is proposed using the nonequilibrium statistical operator method in Renyi statistics.

PACS numbers: 05.20.-y

---

[*] mtok@icmp.lviv.ua

I. INTRODUCTION

During last years, in investigations of complex, self-organizing, fractal structures and various physical phenomena such as subdiffusion, turbulence and chemical reactions, as well as various economical, social and biological systems, Tsallis [1–19], Renyi [20–26, 26–33], Sharma-Mittal [34, 35] statistics as well as superstatistics [36–41] are extensively used along with Gibbs one. The Tsallis entropy is widely applied in various directions of nonextensive statistical mechanics [6, 11–13]. Some examples are the phenomena of subdiffusion [42–45] and turbulence [46–53], and the investigations of transport coefficients in gases and plasma [54, 55], as well as quantum dissipative systems [56]. The problems of the construction of equilibrium thermodynamics in the framework of generalized statistics were discussed in Refs. [57–61]. The energy fluctuations [57], kinetics of nonequilibrium plasma [62], problems of self-gravitating systems [63], the ozone layer [64], universality in non-Debye relaxation [65] and complex systems [66, 67] were investigated within the Tsallis formalism. In Refs. [68–70] Tsallis statistics was applied to description of chemical reactions, in particular, nonlinear equations of reaction-diffusion processes were obtained in Ref. [68]. Despite the wide application of Tsallis entropy as a generalization of Gibbs-Shannon entropy, the Renyi entropy is of great interest as well [71–73]. In particular, in this case it is possible to determine a connection between the index q and the heat capacity of the system [29]. It is important to note the papers [74–76, 78] by Luzzi, Vasconcellos and Ramos where the nonequilibrium statistical operator (NSO) method and the Renyi entropy are applied to description of systems far from equilibrium. In particuar, the nonequilibrium q-dependent Renyi ensemble as well as the generalized distribution functions of bosons and fermions was obtained [74]. A statistical approach for description of fractal physical-chemical systems based on non-Fickian diffusion processes was proposed in Ref. [75]. Therein investigations of anomalous diffusion in fractal-like electrodes in microbatteries were carried out. The experiments on anomalous luminescence at nanometer quantum dots in semiconductor heterostructures were described [76] in such approach. The Refs. [79–81] are devoted to investigation of nonlinear kinetics based on the Kramers, Boltzmann and Fokker-Planck equations within the framework of generalized statistics.

In the present paper, one approach to formulation of an extensive statistical mechanics of nonequilibrium processes [33] is considered, based on the Zubarev NSO method [82–

84] and the maximum entropy principle for the Renyi entropy. This statistical approach is applied to a consistent description of reaction-diffusion processes in the "gas-adsorbate-metal" system. Reaction-diffusion and adsorption-desorption processes on the metal surface are nonlinear, they manifest oscillation character, possess memory effects and are actual in terms of nanostructures formation on surfaces in catalytic phenomena [85–92].

II. RENYI ENTROPY AND NONEQUILIBRIUM STATISTICAL OPERATOR METHOD

The nonequilibrium state of a classical or quantum system of interacting particles is completely described by the nonequilibrium statistical operator $\varrho(x^N;t)$, which satisfies the classical or quantum Liouville equation:

$$\frac{\partial}{\partial t}\varrho(x^N;t) + iL_N\varrho(x^N;t) = 0. \tag{1}$$

iL_N is the Liouville operator of a system.

Within the NSO method framework, we will be looking for solutions of Eq. (1) which are independent of the initial conditions. The solutions will depend explicitly only on the observable quantities

$$\int d\Gamma_N \hat{P}_n \varrho(x^N;t) = \langle\hat{P}_n\rangle^t. \tag{2}$$

The nonequilibrium statistical operator:

$$\varrho(x^N;t) = \varrho_{rel}(x^N;t) - \int_{-\infty}^{t} e^{\varepsilon(t'-t)}T(t,t')\left(1-P_{rel}(t')\right)iL_N\varrho_{rel}(x^N;t')dt',$$

where $T(t,t') = \exp_+\left\{-\int_{t'}^{t}\left(1-P_{rel}(t')\right)iL_N dt'\right\}$ is the evolution operator containing the projection. The relevant statistical operator (distribution function) $\varrho_{rel}(x^N;t)$ will be determined using the maximum entropy principle for the Renyi entropy

$$S^R(\varrho) = \frac{1}{1-q}\ln\int d\Gamma_N\varrho^q(t) \tag{3}$$

at fixed parameters of the reduced description, taking into account the normalization condition.

Relevant statistical operator corresponding to the Renyi entropy maximum:

$$\varrho_{rel}(t) = \frac{1}{Z_R}\left(1 - \frac{q-1}{q}\sum_n F_n(t)\delta\hat{P}_n\right)^{\frac{1}{q-1}}, \tag{4}$$

$$Z_R(t) = \int d\Gamma_N \left(1 - \frac{q-1}{q}\sum_n F_n(t)\delta\hat{P}_n\right)^{\frac{1}{q-1}}. \tag{5}$$

$Z_R(t)$ is the partition function of the relevant statistical operator, $\delta\hat{P}_n = \hat{P}_n - \langle\hat{P}_n\rangle^t$. The Lagrange multipliers $F_n(t)$ are defined from the self-consistency conditions:

$$\langle\hat{P}_n\rangle^t = \langle\hat{P}_n\rangle^t_{rel}, \tag{6}$$

where $\langle\ldots\rangle^t_{rel} = \int d\Gamma_N \ldots \varrho_{rel}(x^N;t)$.

The generalized Kawasaki-Gunton projection operator:

$$P_{rel}(t)\varrho' = \left(\varrho_{rel}(t) - \sum_n \frac{\delta\varrho_{rel}(t)}{\delta\langle\hat{P}_n\rangle^t}\langle\hat{P}_n\rangle^t\right)\int d\Gamma_N\varrho' + \sum_n \frac{\delta\varrho_{rel}(t)}{\delta\langle\hat{P}_n\rangle^t}\int d\Gamma_N\hat{P}_n\varrho'. \tag{7}$$

An action of the operators $P_{rel}(t)iL_N$ on the relevant statistical operator can be presented as follows, by means of generalized projection which now acts on dynamic variables, $P_{rel}(t)iL_N\varrho_{rel}(t) = P_{rel}(t)A(t)\varrho_{rel}(t) = (P(t)A(t))\varrho_{rel}(t)$,

$$P(t)\ldots = \langle\ldots\rangle^t_{rel} + \sum_m \delta\left[\frac{1}{q}\psi^{-1}(t)\left(F_m(t) + \sum_n f^{-1}_{mn}(t)\delta\hat{P}_n\right)\right]\langle\ldots\delta\hat{P}_m\rangle^t_{rel}, \tag{8}$$

Here, $f_{mn}(t)$ is determined by the relations $f_{mn}(t) = \frac{\delta\langle\hat{P}_m\rangle^t}{\delta F_n(t)} = \left(\frac{\delta\langle\hat{P}\rangle^t}{\delta F(t)}\right)_{mn}$,

$$\frac{\delta\langle\hat{P}\rangle^t}{\delta F(t)} = -\left[I - \frac{1}{q}\langle\delta\hat{P}\psi^{-1}(t)\rangle^t_{rel}F\right]^{-1}\frac{1}{q}\langle\delta\hat{P}\psi^{-1}(t)\delta\hat{P}\rangle^t_{rel} = f(t), \tag{9}$$

where I is the unit matrix and $\delta[\ldots] = [\ldots] - \langle[\ldots]\rangle^t_{rel}$. We also use the notation

$$\psi(t) = 1 - \frac{q-1}{q}\sum_n F_n(t)\delta\hat{P}_n. \tag{10}$$

Taking into account that $(1 - P_{rel}(t))iL_N\varrho_{rel}(t) = -\sum_n I_n(t)F_n(t)\varrho_{rel}(t)$, where

$$I_n(t) = (1 - P(t))\frac{1}{q}\psi^{-1}(t)\dot{\hat{P}}_n \tag{11}$$

are the generalized flows, we can now write down an explicit expression for the *nonequilibrium statistical operator*

$$\varrho(x^N;t) = \varrho_{rel}(x^N;t) + \sum_n \int_{-\infty}^t e^{\varepsilon(t'-t)}T(t,t')I_n(t')F_n(t')\varrho_{rel}(x^N;t')dt'. \tag{12}$$

This allows us to obtain the generalized *transport equations* for the reduced-description parameters. They can be presented in the form

$$\frac{\partial}{\partial t}\langle\hat{P}_m\rangle^t = \langle\dot{\hat{P}}_m\rangle^t_{rel} + \sum_n \int_{-\infty}^t e^{\varepsilon(t'-t)}\varphi_{mn}(t,t')F_n(t')dt', \tag{13}$$

with the generalized *transport kernels* (memory functions)

$$\varphi_{mn}(t,t') = \int d\Gamma_N \{\dot{\hat{P}}_m T(t,t') I_n(t') \varrho_{rel}(t')\} \tag{14}$$

which describe the dissipative processes in the system.

III. REACTION-DIFFUSION PROCESSES

Let us start with the Hamiltonian of the system of "gas-adsorbate-metal" in the following form $H = H' + H_{reac}$, $H' = H_a + H_a^{int}$. Here, H_a is the Hamiltonian of the gas subsystem considered according to classical approach; H_a^{int} is the Hamiltonian describing interaction between gas atoms and atoms adsorbed on the metal surface; H_{reac} is the Hamiltonian of interaction for chemical reactions between atoms or molecules adsorbed on the metal surface [85]

$$H_{reac} = \sum_{\bar{a},\bar{b},\bar{a}',\bar{b}'} \left(\langle \bar{a}', \bar{b}' | \Phi_{reac} | \bar{a}, \bar{b} \rangle \hat{q}^{+}_{\bar{a}'} \hat{q}^{+}_{\bar{b}'} \hat{q}_{\bar{a}} \hat{q}_{\bar{b}} + \langle \bar{a}', \bar{b}' | \Phi_{reac} | \bar{a}, \bar{b} \rangle^{*} \hat{q}^{+}_{\bar{a}} \hat{q}^{+}_{\bar{b}} \hat{q}_{\bar{a}'} \hat{q}_{\bar{b}'} \right), \tag{15}$$

where $\langle \bar{a}', \bar{b}' | \Phi_{reac} | \bar{a}, \bar{b} \rangle = \langle \bar{a}, \bar{b} | \Phi_{reac} | \bar{a}', \bar{b}' \rangle$ are the amplitudes of reaction between reagents A and B (supposed to be known from quantum mechanics). We introduce the notation $\bar{a}, \bar{b}$ and $\bar{a}', \bar{b}'$ for state of reagents A, B (atoms or molecules) and for state of atoms in the reaction products AB. Here, $\hat{q}^{+}_{\bar{a}'}$, $\hat{q}^{+}_{\bar{b}'}$, $\hat{q}^{+}_{\bar{a}}$, $\hat{q}^{+}_{\bar{b}}$ and $\hat{q}_{\bar{a}'}$, $\hat{q}_{\bar{b}'}$, $\hat{q}_{\bar{a}}$, $\hat{q}_{\bar{b}}$ are the operators of creation and annihilation of atomic states $\bar{a}', \bar{b}'$ for molecule AB, and $\bar{a}, \bar{b}$ for A and B, respectively.

Parameter of the reduced description are the averaged densities of gas atoms absorbed and not absorbed on the metal surface $\langle \hat{n}_{\bar{a}}(\vec{R}) \rangle^t = Sp\left(\hat{n}_{\bar{a}}(\vec{R})\varrho(t)\right)$, $\langle \hat{n}_{a}(\vec{R}) \rangle^t = Sp\left(\hat{n}_{a}(\vec{R})\varrho(t)\right)$. $\hat{n}_{\bar{a}}(\vec{R})$ is the density operator for gas atoms adsorbed on the metal surface in state ν; $\hat{n}_{\bar{a}}(\vec{R}) = \sum_j^{N_a^{ad}} \hat{\psi}^{+}_{\nu j}(\vec{R}) \hat{\psi}_{\nu j}(\vec{R})$; $\hat{\psi}^{+}_{\nu j}(\vec{R})$, $\hat{\psi}_{\nu j}(\vec{R})$ are the creation and annihilation operators of gas atoms absorbed in state ν on the metal surface which satisfy the Bose-type commutation relations. Since, we do not consider a catalyst surface explicitly in this model, the states ν and μ mean the adsorption centers, where atoms can be located. $\hat{n}_a(\vec{r}) = \sum_j^{N_a^{ad}} \delta(\vec{r} - \vec{r}_j)$ is the microscopic density of gas atoms, $\langle \hat{G}^{\nu\mu}_{\bar{a}\bar{b}}(\vec{R}, \vec{R}') \rangle^t = Sp\left(\hat{G}^{\nu\mu}_{\bar{a}\bar{b}}(\vec{R}, \vec{R}')\varrho(t)\right)$ is the nonequilibrium pair distribution function of atoms or molecules adsorbed on the metal surface, and $\hat{G}^{\nu\mu}_{\bar{a}\bar{b}}(\vec{R}, \vec{R}') = \hat{n}^{\nu}_{\bar{a}}(\vec{R}) \hat{n}^{\mu}_{\bar{b}}(\vec{R}')$.

The relevant statistical operator

$$\varrho_{rel} = \frac{1}{Z_R(t)} \left[1 - \frac{q-1}{q} \beta \left(\delta H(t) - \sum_a \int d\vec{r} \mu_a(r\vec{;}t) \delta \hat{n}_a(r\vec{;}t) \right.\right. \tag{16}$$

$$-\sum_{\bar a}\sum_{\nu}\int d\vec R\mu^{\nu}_{\bar a}(\vec R;t)\delta\hat n^{\nu}_{\bar a}(\vec{R;t})-\sum_{\bar a\bar b}\sum_{\nu\mu}\int d\vec R d\vec R' M^{\nu\mu}_{\bar a\bar b}(\vec R,\vec R';t)\hat G^{\nu\mu}_{\bar a\bar b}(\vec R,\vec R';t)\Bigg)\Bigg]^{\frac{1}{q-1}}.$$

The partition function of the relevant statistical operator

$$Z_R(t)=Sp\Bigg[1-\frac{q-1}{q}\beta\Bigg(\delta H(t)-\sum_a\int d\vec r\mu_a(\vec{r;t})\delta\hat n_a(\vec{r;t}) \tag{17}$$
$$-\sum_{\bar a}\sum_{\nu}\int d\vec R\mu^{\nu}_{\bar a}(\vec R;t)\delta\hat n^{\nu}_{\bar a}(\vec{R;t})-\sum_{\bar a\bar b}\sum_{\nu\mu}\int d\vec R d\vec R' M^{\nu\mu}_{\bar a\bar b}(\vec R,\vec R';t)\hat G^{\nu\mu}_{\bar a\bar b}(\vec R,\vec R';t)\Bigg)\Bigg]^{\frac{1}{q-1}}.$$

$Sp(\ldots)=\prod_\alpha\int\frac{(d\vec r d\vec p)^{N_\alpha}}{N_\alpha!(2\pi\hbar)^{3N_\alpha}}Sp_{(\nu,\xi,\sigma)}(\ldots)$, $N_\alpha=\{N_a,N_{\bar a}\}$, $Sp_{(\nu,\xi,\sigma)}$ means averaged summation over all values of spin and quantum numbers. The parameters $\mu_a(\vec{r;t})$, $\mu^{\nu}_{\bar a}(\vec R;t)$, $M^{\nu\mu}_{\bar a\bar b}(\vec R,\vec R';t)$ should be determined from the corresponding self-consistency conditions

$$\langle\hat n_a(\vec r)\rangle^t=\langle\hat n_a(\vec r)\rangle^t_{rel},\quad\langle\hat n^{\nu}_{\bar a}(\vec R)\rangle^t=\langle\hat n^{\nu}_{\bar a}(\vec R)\rangle^t_{rel},\quad\langle\hat G^{\nu\mu}_{\bar a\bar b}(\vec R,\vec R')\rangle^t=\langle\hat G^{\nu\mu}_{\bar a\bar b}(\vec R,\vec R')\rangle^t_{rel},$$

from where we fined that $\mu_a(\vec{r;t})$ defines a local chemical potential of gas atoms; $\mu^{\nu}_{\bar a}(\vec R;t)$ is a local chemical potential of an atom adsorbed in a state ν on the metal surface. $\delta H(t)=H-\langle H\rangle^t$, $\delta\hat n_a(\vec r;t)=\hat n_a(\vec r)-\langle\hat n_a(\vec r)\rangle^t$, $\delta\hat n^{\nu}_{\bar a}(\vec R;t)=\hat n^{\nu}_{\bar a}(\vec R)-\langle\hat n^{\nu}_{\bar a}(\vec R)\rangle^t$, $\delta\hat G^{\nu\mu}_{\bar a\bar b}(\vec R,\vec R';t)=\hat G^{\nu\mu}_{\bar a\bar b}(\vec R,\vec R')-\langle\hat G^{\nu\mu}_{\bar a\bar b}(\vec R,\vec R')\rangle^t$.

According to (3) and (17), NSO of the system “gas-adsorbate-metal” has the following form

$$\varrho(t)=\varrho(t)_{rel}+\sum_a\int d\vec r\int_{-\infty}^{t}e^{\varepsilon(t'-t)}T(t,t')\left(\int_0^1 d\tau\varrho^{\tau}_{rel}(t')I_a(\vec r;t)\varrho^{1-\tau}_{rel}(t')\beta\mu_a(\vec r;t')\right)dt'$$
$$+\sum_{\bar a}\sum_{\nu}\int d\vec R\int_{-\infty}^{t}e^{\varepsilon(t'-t)}T(t,t')\left(\int_0^1 d\tau\varrho^{\tau}_{rel}(t')I^{\nu}_{\bar a}(\vec R;t)\varrho^{1-\tau}_{rel}(t')\beta\mu^{\nu}_{\bar a}(\vec R;t')\right)dt' \tag{18}$$
$$+\sum_{\bar a,\bar b}\sum_{\nu,\mu}\int d\vec R d\vec R'\int_{-\infty}^{t}e^{\varepsilon(t'-t)}T(t,t')\left(\int_0^1 d\tau\varrho^{\tau}_{rel}(t')I^{\nu\mu}_{G_{\bar a\bar b}}(\vec R,\vec R';t)\varrho^{1-\tau}_{rel}(t')\beta M^{\nu\mu}_{\bar a\bar b}(\vec R,\vec R';t')\right)dt'.$$

$$I_a(\vec r;t)=(1-P(t))\frac{1}{q}\psi^{-1}(t)\dot{\hat n}_a(\vec r),\qquad I^{\nu}_{\bar a}(\vec R;t)=(1-P(t))\frac{1}{q}\psi^{-1}(t)\dot{\hat n}^{\nu}_{\bar a}(\vec R),$$
$$I^{\nu\mu}_{G_{\bar a\bar b}}(\vec R,\vec R';t)=(1-P(t))\frac{1}{q}\psi^{-1}(t)\dot{\hat G}^{\nu\mu}_{\bar a\bar b}(\vec R,\vec R') \tag{19}$$

are the generalized flows describing reaction-diffusion processes. The function $\psi(t)$ is defined by the relation

$$\psi(t)=1-\frac{q-1}{q}\beta\Bigg(\delta H(t)-\sum_a\int d\vec r\mu_a(\vec{r;t})\delta\hat n_a(\vec{r;t}) \tag{20}$$

$$-\sum_{\bar{a}}\sum_{\nu}\int d\vec{R}\mu^{\nu}_{\bar{a}}(\vec{R};t)\delta\hat{n}^{\nu}_{\bar{a}}(\vec{R};t)-\sum_{\bar{a}\bar{b}}\sum_{\nu\mu}\int d\vec{R}d\vec{R}'M^{\nu\mu}_{\bar{a}\bar{b}}(\vec{R},\vec{R}';t)\hat{G}^{\nu\mu}_{\bar{a}\bar{b}}(\vec{R},\vec{R}';t)\Bigg).$$

By means of NSO (19) we obtain the set of self-consistent generalized transport equations for averaged densities of adsorbed and not absorbed atoms along with nonequilibrium pair distribution function of absorbed atoms (or molecules). It has the following form

$$\begin{aligned}\frac{\partial}{\partial t}\langle\hat{n}_a(\vec{r})\rangle^t &= \langle\dot{\hat{n}}_a(\vec{r})\rangle^t_{rel}+\sum_b\int d\vec{r}'\int_{-\infty}^{t}e^{\varepsilon(t'-t)}\varphi_{n_an_b}(\vec{r},\vec{r}';t,t')\beta\mu_b(\vec{r}';t')dt' \\ &+\sum_{\bar{b}}\sum_{\nu'}\int d\vec{R}'\int_{-\infty}^{t}e^{\varepsilon(t'-t)}\varphi^{\nu'}_{n_an_{\bar{b}}}(\vec{r},\vec{R}';t,t')\beta\mu^{\nu'}_{\bar{b}}(\vec{R}';t')dt' \\ &+\sum_{\bar{a}'\bar{b}}\sum_{\nu'\mu'}\int d\vec{R}'\vec{R}''\int_{-\infty}^{t}e^{\varepsilon(t'-t)}\varphi^{\nu'\mu'}_{n_aG_{\bar{a}'\bar{b}}}(\vec{r},\vec{R}',\vec{R}'';t,t')\beta M^{\nu'\mu'}_{\bar{a}'\bar{b}}(\vec{R}',\vec{R}'';t')dt',\end{aligned}\tag{21}$$

$$\begin{aligned}\frac{\partial}{\partial t}\langle\hat{n}^{\nu}_{\bar{a}}(\vec{R})\rangle^t &= \langle\frac{1}{i\hbar}[\hat{n}^{\nu}_{\bar{a}}(\vec{R}),H']\rangle^t_{rel}+\langle\frac{1}{i\hbar}[\hat{n}^{\nu}_{\bar{a}}(\vec{R}),H_{reac}]\rangle^t_{rel} \\ &+\sum_b\int d\vec{r}'\int_{-\infty}^{t}e^{\varepsilon(t'-t)}\varphi^{\nu}_{n_{\bar{a}}n_b}(\vec{R},\vec{r}';t,t')\beta\mu_b(\vec{r}';t')dt' \\ &+\sum_{\bar{b}}\sum_{\nu'}\int d\vec{R}'\int_{-\infty}^{t}e^{\varepsilon(t'-t)}\varphi^{\nu\nu'}_{n_{\bar{a}}n_{\bar{b}}}(\vec{R},\vec{R}';t,t')\beta\mu^{\nu'}_{\bar{b}}(\vec{R}';t')dt' \\ &+\sum_{\bar{a}'\bar{b}}\sum_{\nu'\mu'}\int d\vec{R}'\vec{R}''\int_{-\infty}^{t}e^{\varepsilon(t'-t)}\varphi^{\nu'\mu'}_{n_{\bar{a}}G_{\bar{a}'\bar{b}}}(\vec{r},\vec{R}',\vec{R}'';t,t')\beta M^{\nu'\mu'}_{\bar{a}'\bar{b}}(\vec{R}',\vec{R}'';t')dt',\end{aligned}\tag{22}$$

$$\begin{aligned}\frac{\partial}{\partial t}\langle\hat{G}^{\nu\mu}_{\bar{a}\bar{b}}(\vec{R},\vec{R}')\rangle^t &= \langle\frac{1}{i\hbar}[\hat{G}^{\nu\mu}_{\bar{a}\bar{b}}(\vec{R},\vec{R}'),H']\rangle^t_{rel}+\langle\frac{1}{i\hbar}[\hat{G}^{\nu\mu}_{\bar{a}\bar{b}}(\vec{R},\vec{R}'),H_{reac}]\rangle^t_{rel} \\ &+\sum_{b'}\int d\vec{r}'\int_{-\infty}^{t}e^{\varepsilon(t'-t)}\varphi^{\nu\mu}_{G_{\bar{a}\bar{b}}n_{b'}}(\vec{R},\vec{R}';t,t')\beta\mu_{b'}(\vec{r}';t')dt' \\ &+\sum_{\bar{b}}\sum_{\nu'}\int d\vec{R}''\int_{-\infty}^{t}e^{\varepsilon(t'-t)}\varphi^{\nu\mu\nu'}_{G_{\bar{a}\bar{b}}n_{\bar{b}'}}(\vec{R},\vec{R}',\vec{R}'';t,t')\beta\mu^{\nu'}_{\bar{b}}(\vec{R}';t')dt' \\ &+\sum_{\bar{a}'\bar{b}'}\sum_{\nu'\mu'}\int d\vec{R}''\vec{R}'''\int_{-\infty}^{t}e^{\varepsilon(t'-t)}\varphi^{\nu\mu\nu'\mu'}_{G_{\bar{a}\bar{b}}G_{\bar{a}'\bar{b}}}(\vec{r},\vec{R}',\vec{R}'',\vec{R}''';t,t')\beta M^{\nu'\mu'}_{\bar{a}'\bar{b}'}(\vec{R}'',\vec{R}''';t')dt',\end{aligned}\tag{23}$$

where $\varphi_{n_an_b}$, $\varphi^{\nu\nu'}_{n_{\bar{a}}n_{\bar{b}}}$, $\varphi^{\nu}_{n_an_{\bar{b}}}$, $\varphi^{\nu\mu\nu'\mu'}_{G_{\bar{a}\bar{b}}G_{\bar{a}'\bar{b}}}$ are the generalized transport kernels. The second term in the right-hand side of (23), $\langle\frac{1}{i\hbar}[\hat{n}^{\nu}_{\bar{a}}(\vec{R}),H_{reac}]\rangle^t_{rel}$, defines an averaged value of the operator of rate of the reaction between adsorbed atoms on the metal surface. Transport kernels are built on the generalized flows (20) taking into account contributions of amplitudes of chemical reactions in the flows $I^{\nu}_{\bar{a}}(\vec{R};t')$ and $I^{\nu\mu}_{G_{\bar{a}\bar{b}}}(\vec{R},\vec{R}';t')$, and have the following form

$$\varphi_{BB'}=Sp\left(I_B(t)T(t,t')\int_0^1 d\tau\varrho^{\tau}_{rel}(t')I_{B'}(t')\varrho^{1-\tau}_{rel}(t')\right).\tag{24}$$

In particular, $\varphi_{n_a n_b}(\vec{r},\vec{r}';t,t')$ describes dynamical correlations of diffusive flows of gas atoms and is connected to the inhomogeneous diffusion coefficient of atoms (or molecules) $D_{ab}(\vec{r},\vec{r}';t)$. Similarly, the transport kernel $\varphi^{\nu\nu'}_{n_{\bar{a}} n_{\bar{b}}}(\vec{R},\vec{R}';t,t')$ describes dynamical dissipative correlations of diffusive flows of atoms adsorbed in states ν and ν' on the metal surface and determines inhomogeneous diffusion coefficient of atoms adsorbed on the metal surface $D^{\nu\nu'}_{\bar{a}\bar{b}}(\vec{R},\vec{R}';t)$. Transport kernel $\varphi^{\nu}_{n_{\bar{a}} n_b}(\vec{R},\vec{r}';t,t')$, $\varphi^{\nu'}_{n_a n_{\bar{b}}}(\vec{r},\vec{R}';t,t')$ describes dynamical dissipative correlations between flows of gas atoms and atoms adsorbed on the metal surface, and determines inhomogeneous coefficient of mutual diffusion "gas atom-adsorbed atom" $D^{\nu'}_{a\bar{b}}(\vec{r},\vec{R}';t)$. Investigation of these diffusion coefficients is very important. Transport kernel $\varphi^{\nu\mu}_{G_{\bar{a}\bar{b}}p}$ $(p = n,\bar{n})$ describes dissipative correlations of flows and densities of adsorbed atoms with flows of atoms, molecules and adsorbed atoms. Memory function $\varphi^{\nu\mu\nu'\mu'}_{G_{\bar{a}\bar{b}}G_{\bar{a}'\bar{b}}}(\vec{R},\vec{R}',\vec{R}'',\vec{R}''';t,t')$ describes reaction-diffusion processes between atoms adsorbed on the metal surface. They are higher memory functions with respect to dynamical variables $G^{\nu\mu}_{\bar{a}\bar{b}}$.

IV. CONCLUSIONS

Consequently, we obtain generalized transport equations (22)–(24) for the nonequilibrium averaged densities of adsorbed and not adsorbed atoms of consistent description of the reaction-diffusion processes in the system "gas-adsorbate-metal" within Renyi statistics. At $q = 1$ these equations coincide with the equations of reaction-diffusion processes within Gibbs statistics [85]. As we can see, these equations are nonlinear and spatially inhomogeneous. They can describe strong as well as weak nonequilibrium processes in the system.

---

[1] Tsallis C. Possible Generalization of Boltzmann-Gibbs Statistics. // J. Stat. Phys., 1988., vol. 52, No 1-2., p. 479-489.

[2] Mariz A.M. On the irreversible nature of the Tsallis and Renyi entropies. // Phys. Lett. A, 1992, vol. 165, p.409-411.

[3] Ramshaw J.D. H-theorems for the Tsallis and Renyi entropies.// Phys. Lett. A, 1993, vol. 175, p.169-170.

[4] Ramshaw J.D. Irreversibility and generalized entropies. // Phys. Lett. A, 1993, vol. 175, p.171-172.

[5] Tsallis C. Nonextensive Statistics: Theoretical, Experimental and Computational Evidences and Connections.// Bras. J. Phys., 1999, vol. 29, No 1, p. 1-35

[6] Nonextensive Statistical Mechanics and its Applications; ed. S. Abe and Y. Okamoto. Heidelberg, Spriger-Verlag, 2001., 272 p.

[7] Special issue of Physica A., 2002., Vol. 305, No 1, 2 (edited by G. Kaniadakis, M. Lissia, and A. Rapisarda).

[8] Tsallis C. Nonextensive statistical mechanics: a brief review of its present status.// Anais da Academica Brasileira de Ciencias, 2002, vol. 73, No 3, p. 393-414.

[9] Rajagopal, A. K., Abe S. Implications of Form Invariance to the Structure of Nonextensive Entropies. // Phys. Rev. Lett., 1999., vol. 83., p. 1711-1714.

[10] Abe S., Rajagopal A. K. Microcanonical foundation for systems with power-law distributions. // J. Phys. A., 2000., vol. 33., p. 8733-8739.

[11] Nonextensive Entropy - Interdisciplinary Applications; ed. M. Gell-Mann and C. Tsallis., New York: Oxford Univ. Press, 2004., 440 p.

[12] Tsallis C. Nonadditive entropy and nonextensive statistical mechanics-an overview after 20 years.// Bras. J. Phys., 2009, vol. 99, No 2A, p. 337-356.

[13] Introduction to Nonextensive Statistical Mechanics (Approaching a complex world), Tsallis C. (ed.), Springer, 2009, 389p.

[14] Abe S. Stability of Tsallis entropy and instabilities of Renyi and normalized Tsallis entropies: A basis for q-exponential distributions // Phys. Rev. E., 2002., vol. 66, No 4., p. 046134:1-6.

[15] Abe S., Bagci G. B. Necessity of q-expectation value in nonextensive statistical mechanics. // Phys. Rev. E., 2005., vol. 71, No 1., p. 016139:1-5.

[16] Saguia A., Sarandy M. S. Nonadditive entropy for random quantum spin-S chains.// Phys. Lett. A., 2010., vol. 374., p. 3384-3388.

[17] Lutsko, J. F., Boon J. P., Grosfils P. Is the Tsallis entropy stable? // Europhys. Lett., 2009., vol. 86, p. 40005:1-5.

[18] Tsallis C. From nonlinear statistical mechanics to nonlinear quantum mechanics - concepts

and applications. Proccedings of the Inter. Symp. On Subnuclear Physics: Past, Present and Future., Vatican, 2011, 10p.

[19] Biro T.S., and Molnar E. Non-extensive Statistics, relativistic kinetic theory and fluid dynamics. (EPJ), arXiv: 1205.6079V1, 2012, 10p.

[20] Renyi A. Probability theory. Amsterdam: North-Holland, 1970.

[21] Selected papers by Alfred Renyi, vol. 2, ed. P. Turan., Budapest: Akademiai Kiado, 1976., 662 p.

[22] Beck C., Schlogl F. Thermodynamics of Chaotic Systems. Cambridge, Cambridge Univ. Press, 1993., 308 p.

[23] Klimontovich, Yu. L. Statistical theory of open systems. Moscow, Janus, 1995., 624 p.

[24] Bashkirov A. G., Vityazev A.V. Information entropy and power-law distributions for chaotic systems. // Physica A, 2000, vol. 277, p. 136-145.

[25] Lesche B. Renyi entropies and observables. // Phys. Rev. E., 2004, vol. 70, No 1, p. 017102:1-4.

[26] Jizba P., Arimitsu T. Generalized statistics: yet another generalization. // Physica A, 2004., vol. 340., p. 110-116.

[27] Jizba P., Arimitsu T. The world according to Renyi: thermodenamics of multifracnal systems // Ann. Phys., 2004, vol. 312., p. 17-59.

[28] Masi M. A step beyond Tsallis and Renyi entropies // Phys. Lett. A., 2005, vol. 338., p. 217-224.

[29] Bashkirov A. G., Sukhanov A. D. The distribution function for a subsystem experiencing temperature fluctuations.// JETP., 2002., V. 95, No. 3, . 440.

[30] Bashkirov A. G. On maximum entropy principle, superstatistics, power-law distribution and Renyi parameter // Physica A., 2004., vol. 340., p. 153-162.

[31] Bashkirov A. G. Maximum Renyi Entropy Principle for Systems with Power-Law Hamiltonians. // Phys. Rev. Lett., 2004., vol. 93, No 13., p. 130601:1-4.

[32] Bashkirov A. G. Renyi entropy as a statistical entropy for complex systems. // Theor. Math. Phys., 2006, V. 149, No 2., p. 1559.

[33] Markiv B., Tokarchuk R., Kostrobii P., Tokarchuk M. Nonequilibrium statistical operator method in the Renyi statistics.// Physica A, 2011, vol. 390, p. 785-791.

[34] Sharma B.D., Mittal D.P. // J. Math. Sci., 1975, vol.1, p.28

[35] Akturk E., Bagci G.B., Sever R. Is Sharma-Mittal entropy really a step beyoud Tsallis and

Renyi entropies? arXiv: cond.- mat./ 0703277V1, 2007, 9p.

[36] Beck C. Superstatistics: Theory and Applications. arXiv: cond.- mat./ 0303288V1, 2003, 20p.

[37] Beck C. Superstatistics: Theoretical concept and physical applications. arXiv: cond.- mat./ 0705.3832V1, 2007, 28p.

[38] Abe S., Beck C., Cohen E.G.D. Superstatistics, thermodynamics, and fluctuations

[39] Beck C. Recent developments in superstatistics. arXiv: cond.- mat./ 0811.4363V1, 2009, 7p.

[40] Tsallis C., and Souza A.M.C. Constructing a statistucal mechanics for Beck-Cohen superstatistics. arXiv: cond.- mat./ 0206044V1, 2002, 14p.

[41] Sob'yanin D.N. Generalized of the Beck-Cohen superstatistics. arXiv: cond.- mat./ 1111.6891V1, 2011, 7p.

[42] Essex C., Schulzky C., Franz A., Hoffmann K. H. Tsallis and Renyi entropies in fractional diffusion and entropy production. // Physica A., 2000, vol. 284, p. 299-308.

[43] Tsallis C., Lenzi E.K. Anomalous diffusion: nonlinear fractional Fokker-Planck equation.// Chem. Phys., 2002, vol. 284, p.341-347.

[44] Bologna M., Tsallis C., Grigolini P. Anomalous diffusion associated with nonlinear fractional derivative Fokker-Planck-like equation: Exact time- dependent solutions. arXiv: cond.- mat./ 0003482V1, 2000, 15p.

[45] Boon J.P., Lutsko J.F. Generalized diffusion equation. // Physica A. 2006, vol. 368, p. 55-62.

[46] Beck C., Lewis G.S., Swinney H.I. Measuring non-extensitivity parameters in a turbulent Couette-Taylor flow. arXiv: cond.- mat./ 0010476V1, 2000, 15p.

[47] Arimitsu, T., Arimitsu N. Analysis of fully developed turbulence in terms of Tsallis. // Phys. Rev. E., 2000, vol. 61, No 3, p. 3227-3240.

[48] Ramos F.M., Rosa R.R., Neto C.R., Bolzan M.J.A., Abreu Sa L.D., Camhjs Velho H.F. Non-extensive statistics and three dimensional fully developed turbulence.// Physica A. 2001, vol. 295, p. 250-253.

[49] Arimitsu T., Arimitsu N. Tsallis statistics and turbulence // Chaos, Solutions and Fractals., 2002, vol. 13, p. 479-489.

[50] Arimitsu, T., Arimitsu N. Multifractal analysis of fluid particle accelerations in turbulence.// Physica D., 2004, vol. 193, p. 218-230.

[51] Gotoh T., Kraichnan R.H. Turbulence and Tsallis statistics.// Physica D., 2004, vol. 193, p. 231-244.

[52] Yoshida K. and Arimitsu T. Energy Spectra in Quantum Fluid Turbulence.// J. Low Temp. Phys., 2006, vol. 145, p.219-230.

[53] Arimitsu T., Arimitsu N. and Mouri H. Analyses of turbulence in a wind tunnel by a multifractal theory for probability density functions. arXiv: 1111.4229V3, [physics: flu-dyn], 1012, 17p.

[54] Bezerra, J. R., Silva R., Lima J. A. S. Transport coefficients and nonextensive statistics. // Physica A., 2003, vol. 322, p. 256-266.

[55] Lima, J. A. S., Silva R., Santos J. Plasma oscillations and nonextensive statistics. // Phys. Rev. E., 2000, vol. 61, No 3, p. 3260-3263.

[56] Kirchanov V. S. Using the Renyi entropy to describe quantum dissipative systems in statistical mechanics // Teor. Math. Phys., 2008, vol. 156, No 3, p. 1347-1355.

[57] Feng Z-H., Liu L-Y. Energy fluctuation and correlation in Tsallis statistics. // Physica A, 2010, vol. 389, p. 237-241.

[58] Boon J. P., Lutsko J. F. Is nonextensive statistics applicable to continuous Hamiltonian systems? arXiv:1003.3592v1 [cond-mat. stat-mech], 2010, 6 p.

[59] Du J. L. A new form of Tsallis distribution based on the probabilistically independent postulate. // Chin. Phys. B, 2010, vol. 19, p. 070501:1-11.

[60] Keshavarzi E. M. Sabzehzari M., Eliasi M. Quantum vibrational partition in the non-extensive Tsallis framework.// Physica A, 2010, vol. 389, p. 2733-2738.

[61] Hasegawa H. The interpolation approach to nonextensive quantum statistics. // Physica A, 2010, vol. 389, p. 2358-2375.

[62] Du J. L. Nonextensivity in nonequilibrium plasma systems with Coulombian long-range interactions. //Phys. Lett. A, 2004, vol. 329, p. 262-267.

[63] Du J. L. Jeans' criterion in nonextensive statistical mechanics.// Physica A, 2004, vol. 335, p. 107-114.

[64] Ferri G. L., Ferri G. L., Savio M. F. R., Plastino A. Tsallis' q-triplet and the ozone layer. // Physica A, 2010, vol. 389, p. 1829-1833.

[65] Brouers F., Sotolongo-Costa O., Weron K. Burr, Levy, Tsallis. // Physica A, 2004, vol. 344, p. 409-416.

[66] Yulmetyev R. M., Gafarov F. M., Yulmetyeva D. G., Emeljanova N. A. Intensity approximation of random fluctuation in complex systems. // Physica A, 2002, vol. 303, p. 427-438.

[67] Yulmetyev R. M., Emeljanova N. A., Gafarov F. M. Dynamical Shannon entropy and information Tsallis entropy in complex systems. // Physica A, 2004, vol. 341, p. 649-676.

[68] Plastino A.R., Casas M., Plastino A. A nonextensive maximum entropy approach to a family of nonlinear reaction-diffusion equations. // Physica, 2000, vol. 280, p. 289-303.

[69] Niven R.K. q-Exponential Structure of Arbitrary-Order Reaction Kinetics. Shorter Communication., 2005, 17p.

[70] Bagci G.B. Nonextensive Reaction Rate. arXiv: 0705.2050V1, [cond-mat. stat-mech], 2007, 6p.

[71] Parvan A. S., Biro T. S. Extensive Renyi statistics from non-extensive entropy.// Phys. Lett. A, 2005, vol. 340, p. 375-387.

[72] Parvan A. S., Biro T. S. Renyi statistics in equlibrium statistical mechanics.// Phys. Lett. A (in press).

[73] Figueiredo A., Amato M. A., Filho M. T. R. On the statistical interpretation of generalized entropies . // Physica A. - 2006. - Vol. 367. - P. 191-206.

[74] Luzzi R., Vasconcellos A.R., and Ramos J.G. Competing styles of statistical mechanics: I. Systematization and Clarification in a General Theory. arXiv.org/abs/cond-mat/0306217, 2004, 44p.

[75] Luzzi R., Vasconcellos A.R., and Ramos J.G. Competing styles of statistical mechanics: II. Comparision of Theory and Experiment and Further Illustrations. arXiv.org/abs/cond-mat/0306247, 2004, 40p.

[76] Vasconcellos A.R., and Ramos J.G., Gorenstein A., Kleinke M.U., Souza Cruz T.G., Luzzi R. Statistical Approach to Fractal-Structured Physico-Chemical Systems: Analysis of non-Fician diffusion. arXiv: cond-mat/0412131V1 [cond-mat. stat-mech], 2004, 28p.

[77] Luzzi R., Vasconcellos A.R., and Ramos J.G. Entropy: Mystery and Controversy. Plethora of Informatinal Entropies and Unconventional Statistics. arXiv.org/abs/cond-mat/0307235, 2004, p.

[78] Luzzi R., Vasconcellos A.R., and Ramos J.G. On nonextensive thermo-statistics: systematization, clarification of scope and interpretation, and illustrations. arXiv: cond-mat/0412233V1 [cond-mat. stat-mech], 2004, 33p.

[79] Kaniadakis G. Non-linear kinetics underlying generalized statistics. // Physica, vol. 296, p.405-425.

[80] Borland L. Microscopic dynamics of the nonlinear Fokker-Planck equation: A phenomenological model. // Phys. Rev. E, 1998, vol.57, No 6, p.6634-6642.

[81] Frank T.D. Nonlinear Fokker-Planck Equations. Fundamentals and Applications. Springer, Berline Heid, New York, 2004, 413p.

[82] Zubarev D. N. Nonequilibrium Statistical Thermodynamics. New-York: Consultant Bureau, 1974. - 489 p.

[83] Zubarev D. N., Morozov V. G., Repke G. Statistical Mechanics of Nonequilibrium Processes: 1. Basic Concepts, Kinetic Theory. Berlin: Alademie Verlag, 1996, 270 p.

[84] Zubarev D. N., Morozov V. G., Repke G. Statistical Mechanics of Nonequilibrium Processes. Moscow, Fizmatlit, 2002, 432 p. (in Russian).

[85] Kostrobii P.P., Tokarchuk M.V., Markovych B.M., Ignatyuk V.V., Gnativ B.V. Reaction-diffusion processes in the“metal-gas” systems. Publishing House of Lviv Polytechnic National University, Lviv, 2009 (in Ukrainian).

[86] Ala-Nissila T., Ferrando R. and Ying S.C.// Adv. Phys., 2002, vol. 51(3), p.949

[87] Ignatyuk V.V. Kinetic equation approach to the description of quantum surface diffusion: Non-Markovian effects versus jump dynamics. // Phys. Rev. E., 2009, V.80, P.041133:1-13.

[88] Ignatyuk V.V. Coherence, decoherence, and memory effects in the problems of quantum surface diffusion. // Phys. Rev. E., 2011, V.84, P.021111:1-13.

[89] Ignatyuk V.V. A temperature behavior of the frustruted translational mode of adsorbate and the nature of the ”adsorbate-substrate” interaction.// J Chem. Phys., 2012, vol. 136, p. 184104-11

[90] Galenko P.K., Kharchenko D., Lysenko I.Stochastic generalization for a hyperbolic model of spinodal decomposition. // Physica A. –2010. –V.389. –P.3443–3455.

[91] Ghosh P., Sen S., Ray D. S. Spatial periodicity induced by a chemical wave train. // Phys. Rev. E., 2010, V.80, P. 017101:1-4.

[92] Kharchenko V.O., Kharchenko D.O., Kokhan S.V., Vernigora I.V., Yanovsky V.V. Properties of nano-islands formation in nonequilibrium reaction-diffusion systems with memory effects.arXiv: 1206.0174V1, [cond-mat.stat-mech], 2012, 12p.